

The Biodiversity and Farmer Practices in Different Climate and

Their Effect on The Infestation Rate of Coffee Berry Borer

Marich Nur MAQSALINA^{1,2}, Leïla Bagny BEILHE³, Wagianto WAGIANTO⁴, Purnama HIDAYAT⁵

^{1,2} IPB University and University of Montpellier; ³ PHIM, Univ. Montpellier, CIRAD, INRAE, Institut Agro, IRD, Montpellier, France; ⁴ PT Indo Cafco, North Sumatera, Indonesia; ⁵ Departement of Plant Protection, IPB University, Bogor, Indonesia



Introduction

The coffee berry borer (CBB), *Hypothenemus hampei* is a major coffee pest that attacks and spreads to all coffee producing countries, including Indonesia [1] causing important yield and economic losses [2]. CBB incidence can be modulated by farmers' practices, and by biotic and abiotic factors.

In this study, we propose to investigate the effects of all these factors on CBB infestation rate in two regions of Indonesia, Simalungun Regency in North Sumatera and Bandung Regency in West Java. Both regions are mainly composed of Arabica coffee plantations [3]. The North Sumatera region is submitted to an equatorial climate which has a bimodal type of rainfall whereas the West Java region is a typical tropical humid climate with only one peak of rain.

Methods

Twenty study plots were selected per region at an altitude around 1200m, with two levels of practices regarding CBB control (10 farms with control strategy and 10 farms without control) and based on the number of associated trees species that are present.

In each farm, all coffee berries (attacked or not by CBB) were counted on 120 branches on 30 randomly selected trees during five months (from January to May 2023). Infestation rate of CBB was calculated monthly. Farmers used to associate more trees species to coffee plantation in North Sumatera (6 max) than in West Java (14 max).




Figure 1: The percentage of dynamics of CBB infestation rates in two areas




Figure 2: The percentage of infestation rates of CBB per 5 months based on area




Figure 3: The percentage of infestation rates of CBB per 5 months based on CBB control

Results

Global incidence of CBB is higher in North Sumatera than in West Java whatever the level of associated trees species. The farmer practices affect on the infestation rate of CBB. However, the effect of climate on this dynamics is under investigation. Moreover, the effect of the number of associated trees species on CBB incidence is still unclear. We assumed that the effect of associated biodiversity on CBB is linked to microclimate modification and additional data are needed to confirm that. Our result demonstrated that control strategies implemented by farmers are efficient to reduce CBB infestation.

Conclusion & Perspectives

Results obtained in this study have never been published for these regions. Based on these results we will adapt an existing agent-based model developed in Costa Rica on CBB infestation rate dynamics. This model will then be presented to Indonesian farmers to support implementation of efficient control strategies that are adapted to local context.

References

- [1] R. Harni *et al.*, *Coffee Pest and Disease Control Technology*. Jakarta: Indonesian Agency for Agricultural Research and Development (IAARD) Press, 2015.; [2] J. R. Saragih, "Socioeconomic and Ecological Dimension of Certified and Conventional Arabica Coffee Production in North Sumatera, Indonesia," *Asian J. Agric. Rural Dev.*, vol. 3, no. 3, pp. 93–107, 2013, [Online]. Available: [http://www.aessweb.com/pdf-files/2-186-AJAR-3\(3\)2013-93-107.pdf](http://www.aessweb.com/pdf-files/2-186-AJAR-3(3)2013-93-107.pdf); [3] A. Bustillo, *et al.*, *Manejo Integrado de la Broca del Cafe*. Colombia: Cenicafe, 1988.